

## Detachment of crash helmets during motorcycle accidents

Since 1973 motorcyclists have been required to wear a properly fastened, approved helmet when on the road. I report four cases in which such helmets became detached during an accident. In three cases fatal injuries resulted.

### Case reports


**Case 1**—An 18 year old girl riding pillion on a motorcycle was thrown from the machine after a collision with an oncoming vehicle. Her helmet was seen to fly off before impact and was found later with intact strap and fixing bolts. She was unconscious from impact and suffered diffuse brain injury. She died five days later.

**Case 2**—A 21 year old racing motorcyclist crashed, and during the course of the accident I witnessed his helmet become detached from his head; fortunately he was not injured. The helmet was found with the strap correctly fastened and the fittings intact. Later examination of the rider with the helmet on showed that an apparently well fitting helmet could be easily lifted over the back of his head.

**Cases 3 and 4**—Two racing motorcyclists, aged 23 and 29, collided and were thrown from their machines. A film of the accident showed both helmets clearly coming off over the backs of the riders' heads. Both riders suffered fatal diffuse brain injuries.

### Comment

Current motorcycle helmets are designed to reduce the severity of an injury by three mechanisms: firstly, by presenting a hard surface to prevent direct blows to the skull; secondly, by having a deformable polystyrene liner within the shell to dissipate deceleration forces; and, thirdly, by presenting a smooth surface to the road to prevent rotation of the head and neck. The helmet is strapped to the head with woven or leather straps that are fixed to the helmet by rivets. Failure of the shell, lining, or strap may lead to failure of the helmet itself, and uniformity of strength is controlled by the British Standards Institution.<sup>1,2</sup>



Manner in which helmet is detached at high speed, showing (left) normal position and (right) helmet pivoting forward on strap during accident.

All patients reported on here were wearing correctly fastened, full face helmets. Witnesses clearly saw the helmets fly off, and there was no question of the helmets having been removed after the patient had come to rest. There was no major damage to the shell or lining of the helmets recovered in the three fatal cases. One patient (case 2) was not injured, and the mechanism of the helmet coming off could be seen. The fixing bolts of the chin strap were near the base and the front of the helmet. This allowed the helmet to pivot forward on the strap, the pivot point lying anterior to the angle of the mandible (figure). This is more likely to happen if the rider has a flattened occiput. When I tested this on a random group of 14 motorcyclists with apparently well fitting, comfortable, and properly fastened helmets I was able to remove four of the helmets by pushing from behind.

Many motorcyclists may possibly be at risk of losing their helmets in accidents. Modification of the helmet by moving the strap mounting bolts further back and higher up the helmet would alter the pivot point, making it harder for the helmet to come off. Alternatively, the customer and dealer selling helmets should be aware of the prob-

lem. A simple test of somebody pushing the helmet upwards from behind while it is on the customer's head should be performed. If the helmet can reach the vertex or beyond the customer should refuse to buy it and the dealer to sell it. The four patients described here were wearing four different makes of helmet, indicating that the problem is widespread. The manufacturer of the helmet in case 2 has already modified the product. It is to be hoped that other manufacturers will take a similarly responsible attitude.

I thank Mrs Irene Prentice for her artistic help.

<sup>1</sup> British Standards Institution. *Specification for protective helmets for vehicle users*. BS5361. London: British Standards Institution, 1976.

<sup>2</sup> British Standards Institution. *Specification for protective helmets for vehicle users (high protection)*. BS2495. London: British Standards Institution, 1977.

(Accepted 17 November 1983)

Central Middlesex Hospital, London NW10

P G RICHARDS, FRCS, neurosurgical registrar

Correspondence to: Mr P G Richards, Gough-Cooper Department of Neurological Surgery, Institute of Neurology, London WC1.

## Role of anaerobes in cerebral abscesses of sinus origin

Cerebral abscesses of sinus origin have been attributed to microaerophilic streptococci and treated with penicillin.<sup>1</sup> We report three consecutive cases seen in our department, each of which grew a mixed aerobic and anaerobic microbacterial flora, including penicillin resistant organisms.

### Case reports

**Case 1**—A 20 year old man developed right ethmoiditis associated with proptosis and periorbital cellulitis. Right ethmoidectomy and drainage of the frontal sinus were performed. During treatment with benzylpenicillin and chloramphenicol he developed left frontal sinusitis. On exploration the posterior frontal sinus wall was found to be deficient. Metronidazole was added to the regimen. The patient deteriorated and a right frontal lobe abscess was drained, yielding *Streptococcus milleri* Lancefield group C, *Haemophilus aphrophilus*, and *Bacteroides melaninogenicus*, resistant to penicillin. Further deterioration necessitated a partial right frontal lobectomy, from which he made a good recovery.

**Case 2**—A 15 year old boy with frontal sinusitis was treated with amoxycillin. He developed a right hemiparesis and focal fits. An extradural left frontal abscess and left pansinusitis were confirmed by computed tomography and evacuated. During treatment with benzylpenicillin and sulphadimidine the patient deteriorated and a left subdural abscess was aspirated. Anaerobic streptococci and *Str milleri* Lancefield group F were isolated from the abscess and pus fluoresced red under ultraviolet light, suggesting the presence of *B melaninogenicus*. Metronidazole was added to the regimen and the patient made a full recovery.

**Case 3**—A 24 year old man with pansinusitis treated by oral penicillin was admitted with left sided focal fits and hemiparesis. Right subdural empyema and pansinusitis were confirmed by computed tomography and pus evacuated from these sites yielded *Fusobacterium* spp, anaerobic streptococci, and *H influenzae*. The patient had begun benzylpenicillin, gentamicin, and metronidazole on admission and made a full recovery.

### Comment

It is well recognised that cerebral abscesses of otitic aetiology frequently yield a mixed aerobic and anaerobic microbacterial flora, often including *Bacteroides* spp. In contrast, cerebral abscesses of sinusitic origin have been attributed primarily to "penicillin sensitive streptococci" and it has been asserted that chemotherapy should be aimed at these organisms.<sup>1</sup>

A recent study has shown that potentially pathogenic anaerobes are usually present in the normal paranasal sinus, with *Peptostreptococcus* spp and *Bacteroides* spp most commonly cultured.<sup>2</sup> When purulent secretion is present the oxygen tension of the maxillary sinus is zero, favouring the proliferation of these organisms. Other studies have shown that similar anaerobes are frequently present in both acute and chronic maxillary sinusitis.<sup>3</sup>

In each of our cases a mixed aerobic and anaerobic flora was